


Early Photoreceptor Apoptosis and Microglial Response in Light-Damaged Mouse Retina

Hanieh Shojaee¹, Nourani M.R.^{2,*} 

1. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

2. Tissue Engineering and Regenerative Medicine, New Health Technologies Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran.

*. Corresponding author: Mohammad reza Nourani, r.nourani@yahoo.com



Hanieh Shojaee

Article type:

Original Article

Received: 26 Nov. 2025

Revised: 12 Dec. 2025

Accepted: 20 Dec. 2025

ePublished: 25 Dec. 2025

Abstract

Background: The retina, as part of the central nervous system, has limited capacity for self-repair following injury. Light-induced retinopathy serves as a valuable model for studying retinal degeneration, but the early events following photic injury remain incompletely characterized.

Objective: This study investigated the temporal and spatial relationship between photoreceptor apoptosis and macrophage response during the first three days after light-induced retinal damage.

Methods: Adult BALB/c mice were exposed to intense cool white fluorescent light (3,500-7,000 lux) for 7 hours and sacrificed at 1, 2, and 3 days post-injury. Retinal sections were examined using toluidine blue staining for morphology, TUNEL assay for apoptotic cells, and F4/80 immunohistochemistry for macrophages/microglia.

Results: Progressive outer nuclear layer (ONL) thinning was observed, with near-complete photoreceptor loss by day three. TUNEL-positive apoptotic cells appeared exclusively in the ONL, peaking at day two and declining by day three. F4/80-positive cells transformed from ramified to amoeboid morphology, increased in number, and migrated to the ONL, with peak accumulation at days two to three.

Conclusion: Photoreceptor apoptosis peaks at 48 hours post-injury, closely followed by microglial activation and recruitment to the site of damage. This temporal sequence confirms a direct relationship between cell death and inflammatory response, identifying the first 48 hours as a critical therapeutic window for intervention in retinal degenerative diseases.

Keyword: Photoreceptor Apoptosis, Microglial, Light-Damaged, Mouse Retina

Introduction

The retina, as the light-sensitive tissue at the back of the eye, is particularly vulnerable to photochemical damage due to its high metabolic rate, abundant polyunsaturated fatty acids, and constant exposure to light [1, 2]. Excessive or prolonged exposure to light can trigger a cascade of degenerative events in photore-

ceptor cells, a process known as light-induced retinopathy. This pathological phenomenon is not only a significant concern in the context of occupational or environmental light overexposure but also serves as a valuable experimental model for understanding the mechanisms underlying prevalent retinal degenerative diseases such as Age-Related Macular Degeneration (AMD) and Retinitis Pigmentosa [3, 4]. In these

conditions, oxidative stress and subsequent cell death are key drivers of vision loss, making the light-damage model a powerful tool for investigating these fundamental pathological processes. Histologically, the mammalian retina is a highly organized, multi-layered structure. The outer nuclear layer (ONL) contains the cell bodies of photoreceptors rods and cones which are the primary victims of light-induced damage [5]. Beneath this lies the inner nuclear layer (INL), housing the cell bodies of bipolar, horizontal, and amacrine cells, while the ganglion cell layer (GCL) contains the cell bodies of retinal ganglion cells whose axons form the optic nerve. The photore-

ceptor outer segments interdigitate with the retinal pigment epithelium (RPE), a monolayer of cells critical for phagocytosis of shed outer segments, recycling of visual cycle components, and maintaining the blood-retina barrier [6]. The integrity of these layers, particularly the ONL and RPE, is essential for visual function, and their disruption is a hallmark of retinal degeneration. This highly structured architecture allows for precise histological localization of pathological events, such as apoptotic nuclei in the ONL or infiltrating immune cells within specific layers (Fig 1).




Figure 1: Histological structure of retina

A central feature of retinal degeneration, regardless of the initial insult, is the programmed death of photoreceptors, primarily via apoptosis [7, 8]. This process has been extensively documented using techniques like TUNEL (Terminal deoxynucleotidyl transferase dUTP nick end labeling), which specifically labels fragmented DNA in apoptotic cells [9, 10]. Following this cell death, a robust inflammatory response is triggered. Resident microglia, the im-

mune cells of the central nervous system, become activated and, along with infiltrating macrophages from the circulation, migrate to the site of injury [11, 12]. These cells, collectively identified by markers such as F4/80 in rodents, play a dual role: they are essential for clearing cellular debris, but they can also exacerbate tissue damage by releasing pro-inflammatory cytokines [13, 14]. The delicate balance between these protective and destructive roles

is a critical determinant of the final outcome for retinal tissue. While the individual phenomena of apoptosis and macrophage infiltration in the damaged retina are well-established, the precise temporal and spatial relationship between them in the very first few days following injury remains incompletely characterized [15]. Understanding this early phase is crucial, as it represents a potential window for therapeutic intervention. Most studies have focused on later time points where degeneration is fully established, leaving a gap in our knowledge of the initiating events and the immediate cellular dialogue between dying neurons and responding immune cells.

Therefore, this study was designed to investigate the acute phase of light-induced retinal damage in a mouse model. Our primary objective was to characterize the time course of apoptotic cell death and the concurrent macrophage response within the first three days post-lesion. Using TUNEL assay to label apoptotic cells and immunohistochemistry for the macrophage marker F4/80, we aimed to provide a detailed histological timeline of these key degenerative and inflammatory events. We hypothesized that the peak of macrophage infiltration would closely follow the peak of apoptosis, suggesting a direct recruitment mechanism in this early stage of pathology. The findings from this study will contribute to a better understanding of the initial pathophysiological processes in retinal degeneration and may help identify targets for early therapeutic strategies.

Materials and Methods

Animals

Eight-week-old male BALB/c mice were used in this study. All animals were housed in ventilated cages under controlled environmental conditions (22°C, 12:12-hour light/dark cycle) with food and water available ad libitum. For tissue collection, mice were deeply anesthetized with an intraperitoneal injection of pentobarbital (50 mg/kg body weight) and subsequently perfused transcardially with 4% paraformaldehyde in 0.1 M phosphate buffer (PB), pH 7.2. Following enucleation, eyeballs were post-fixed in the same fixative for 2 hours at 4°C and prepared for histopathological and immunohistochemical analysis

using cryosectioning. All experimental procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of Laboratory Animal Sciences Center.

Light-Induced Retinal Damage Model

Photic injury was induced according to the protocol previously described [16]. Thirteen-week-old male BALB/c mice were randomly assigned to two experimental groups: light-exposed (LE) and control.

Control mice were dark-adapted for 24 hours and then exposed to standard room illumination (approximately 200 lux) for 6 hours prior to euthanasia. Light-exposed mice were similarly dark-adapted for 24 hours and subsequently exposed to intense cool white fluorescent light at an intensity ranging from 3,500 to 7,000 lux for 7 hours, beginning at 2:00 a.m. and ending at 9:00 a.m. [17]. Following photic injury, LE mice were returned to the normal 12:12-hour light/dark cycle until euthanasia.

Animals were sacrificed at 6 hours, 1 day, 2 days, and 3 days post-exposure. At each time point, a minimum of three mice were used for tissue collection. Age-matched control mice (n=3) were also included.

Tissue Preparation

Following transcardial perfusion, eyeballs were enucleated and post-fixed in 4% paraformaldehyde in 0.1 M PB (pH 7.2) for 2 hours at 4°C. The tissues were then cryoprotected by immersion in graded sucrose solutions (10%, 20%, and 30%) in 0.1 M PB at 4°C until sinking. Specimens were embedded in optimal cutting temperature (OCT) compound and rapidly frozen. Serial cryosections were cut at a thickness of 10–12 μ m in the sagittal plane through the optic nerve head, collected on gelatin-coated slides, and stored at –20°C until further processing.

TUNEL Assay for Apoptotic Cell Detection

Apoptotic cells in retinal sections were identified using the terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling (TUNEL) technique [9]. Cryosections were processed according to the manufacturer's instructions using the In Situ Cell Death Detection Kit (Roche Diagnostics). Briefly,

sections were permeabilized with 0.1% Triton X-100 in 0.1% sodium citrate for 2 minutes on ice, followed by incubation with the TUNEL reaction mixture for 60 minutes at 37°C in a humidified chamber. Negative controls were prepared by omitting the terminal deoxynucleotidyl transferase enzyme. Sections were counterstained with 4',6-diamidino-2-phenylindole (DAPI) to visualize all cell nuclei. TUNEL-positive cells were examined under a fluorescence microscope.

Immunohistochemistry for Macrophages

For immunohistochemical detection of macrophages, cryosections were first permeabilized with 0.1% Triton X-100 in 0.1 M phosphate-buffered saline (PBS) for 10 minutes. Endogenous peroxidase activity was quenched by incubation with 0.3% hydrogen peroxide (H₂O₂) in methanol for 20 minutes at room temperature. To block non-specific antibody binding, sections were incubated with 1% bovine serum albumin (BSA) in 0.1 M PBS for 1 hour at room temperature [18].

Sections were then incubated overnight at 4°C with a primary antibody against F4/80, a specific marker for murine macrophages (rat anti-mouse F4/80, 1:500 dilution; AbD Serotec, Oxford, UK) [19]. After thorough washing with PBS, sections were incubated with appropriate biotinylated secondary antibody (goat anti-rat IgG, 1:200; Vector Laboratories) for 1 hour at room temperature. Immunoreactivity was visualized using the Vectastain Standard HRP Kit (Vector Laboratories, Burlingame, CA, USA) with 3,3'-diaminobenzidine (DAB, 0.05%) as the chromogen. The reaction was monitored under a microscope and stopped by rinsing with distilled water. Sections were dehydrated through graded ethanol, cleared in xylene, and mounted with Permount. Negative control sections were processed by omitting the primary antibody.

Microscopy and Image Analysis

Stained sections were examined for TUNEL-positive and F4/80-positive cells using an Olympus BX51 light/fluorescence microscope. Images were captured using a DP72 digital camera (Olympus,

Japan) and analyzed with image analysis software (ImageJ, NIH).

Results

As the retina is part of the central nervous system (CNS), it possesses a very limited capacity for self-repair. Following an injury caused by light exposure, the retina undergoes rapid degeneration, with the most critical damage occurring in the outer nuclear layer (ONL), which contains the photoreceptor cell bodies [5]. At 1, 2, and 3 days post-injury, this layer became progressively disorganized, and its thickness decreased by more than half, as revealed by toluidine blue staining in Figure 2 [20].

In the intact retina, F4/80-immunopositive cells were sparsely distributed throughout the retinal layers, displaying a ramified morphology with arborized, thin processes characteristic of a resting, surveillance state [19]. Following photic injury, these cells increased in number and transitioned to a round, amoeboid morphology, signifying their shift to an activated, phagocytic state [11]. This morphological change coincided with their relocation to the ONL. As degeneration proceeded, the F4/80-positive cells continued to proliferate, becoming larger and more rounded, and ultimately concentrated within the areas of the outer retina undergoing active remodeling [14].

To assess apoptosis following photic injury, TUNEL staining was performed. In the intact retina, no TUNEL-positive cells were observed throughout the entire retinal tissue. Following photic injury, the number of apoptotic cells gradually increased [10]. These cells were predominantly localized within the ONL. The peak frequency of TUNEL-positive cells was observed at 2 days post-injury [8]. Subsequently, as the thickness of the ONL decreased, the number of TUNEL-positive cells declined. This temporal pattern indicates that the majority of photoreceptor cell death occurs around 2 days following photic injury [7].




Figure 2: Histological and immunohistochemical analysis of retinal degeneration after photic injury

Retinal sections from intact and injured mice were stained with toluidine blue to visualize general morphology (A-D), with an antibody against F4/80 to detect microglia (E-H), and with TUNEL to identify apoptotic photoreceptors (I-L). (A-D) Toluidine blue staining. (A) Intact retina showing the distinct retinal layers: ganglion cell layer (GCL), inner nuclear layer (INL), outer nuclear layer (ONL), and retinal pigment epithelium (RPE). (B) Retina at 1 day post-injury, showing early disorganization of the ONL. (C) Retina at 2 days post-injury, with significant thinning of the ONL. (D) Retina at 3 days post-injury, where most photoreceptor nuclei have disappeared. (E-H) F4/80 immunohistochemistry. (E) In the intact retina, rare F4/80-positive microglia display a ramified morphology with thin, arborized processes. (F) At 1 day post-injury, microglia increase in number and begin to migrate to the ONL. (G) At 2 days post-injury, microglia are predominantly round/amoeboid in shape and localized within the ONL. (H) At 3 days post-injury, numerous rounded microglia are concentrated in the area of active ONL remodeling. (I-L) TUNEL assay. (I) The intact retina shows no TUNEL-positive cells [9]. (J) At 1 day post-injury, scattered TUNEL-

Discussion

The present study was designed to investigate the acute phase of light-induced retinal damage in a mouse model, focusing on the temporal and spatial relationship between photoreceptor apoptosis and the subsequent macrophage response during the first three days post-injury. Using toluidine blue staining,

TUNEL assay, and F4/80 immunohistochemistry, we aimed to provide a comprehensive histological timeline of degenerative and inflammatory events following light exposure [15]. Our results confirm the hypothesis that the peak of macrophage/microglial accumulation in the outer nuclear layer (ONL) closely follows the peak of photoreceptor apoptosis, supporting the concept that dying photo-

receptors release chemotactic signals that recruit immune cells to the site of injury [12].

Toluidine blue staining revealed progressive and severe degeneration of the ONL following light exposure. In intact retinas, the characteristic multi-layered architecture was clearly discernible. However, by day one post-injury, disorganization of the ONL was evident. At day two, significant thinning of this layer was observed, and by day three post-injury, the majority of photoreceptor nuclei had disappeared, with the ONL thickness reduced by more than half [20]. These findings align with previous reports demonstrating that photoreceptors are the primary targets of light-induced damage due to their high metabolic activity and susceptibility to oxidative stress [1, 2]. The rapid progression within 72 hours corroborates earlier studies reporting that the timing of photoreceptor loss depends on light intensity and genetic background [17]. The near-complete ONL loss by day three underscores the acute nature of this model and highlights the narrow window for neuroprotective intervention [3].

TUNEL staining confirmed that photoreceptor loss occurred via apoptosis. No TUNEL-positive cells were detected in intact retinas. At day one post-injury, scattered TUNEL-positive cells appeared exclusively within the ONL. This number increased progressively, reaching a peak at day two post-injury, then declined by day three as the ONL thinned [8]. This temporal pattern strongly agrees with previous studies demonstrating that light-induced photoreceptor apoptosis in BALB/c mice peaks between 24 and 48 hours, with maximal DNA fragmentation observed at 48 hours [7, 10]. The decline by day three reflects that most susceptible photoreceptors had already undergone cell death and been cleared, consistent with the concurrent ONL thinning observed in our study [9]. The exclusive ONL localization of TUNEL-positive cells reinforces that photoreceptor apoptosis is the primary pathological mechanism driving early retinal degeneration, validating the light-damage model for studying photoreceptor cell death mechanisms relevant to AMD and retinitis pigmentosa [4, 6].

In parallel, we characterized F4/80-immunopositive cells representing resident microglia and infiltrating macrophages. In intact retinas, these cells were sparsely distributed across all retinal layers and exhibited a ramified morphology with thin, arborized processes, characteristic of resting microglia in their surveillance state [19]. Following light exposure, a dramatic transformation occurred. By day one post-injury, these cells had increased in number and begun to migrate toward the ONL. At day two, they had adopted a round, amoeboid morphology—a hallmark of microglial activation and transition to a phagocytic phenotype and were predominantly localized within the degenerating ONL [14]. By day three, numerous rounded F4/80-positive cells were concentrated in the areas of active ONL remodeling [13]. This progression from ramified to amoeboid morphology, coupled with redistribution to the injury site, is consistent with established patterns of microglial activation following neuronal damage [11, 18]. Our findings align with studies demonstrating microglial activation within 24 hours of light-induced injury and migration to the ONL to phagocytose apoptotic debris, with activated microglial accumulation peaking between days two and three [12, 14].

A central objective was examining the temporal relationship between these events. Our results demonstrate a clear sequence: photoreceptor apoptosis begins within 24 hours, peaks at 48 hours, and declines by 72 hours. Concurrently, microglial activation and migration become evident by day one, peak between days two and three, and persist through day three in areas of active remodeling [15]. This temporal cascade confirms our hypothesis: the peak of microglial accumulation slightly lags behind the peak of apoptosis, consistent with the time required for microglia to sense damage signals, become activated, and migrate [14]. The close spatial relationship between TUNEL-positive and F4/80-positive cells within the ONL further supports that microglia directly respond to apoptotic photoreceptors, likely through "find-me" signals such as ATP and fractalkine [13]. Once recruited, microglia fulfill their essential phagocytic role, clearing cellular debris to prevent sec-

ondary necrosis and inflammation [18]. However, activated microglia also secrete pro-inflammatory cytokines (TNF- α , IL-1 β) that may exacerbate tissue damage [11]. The persistence of activated microglia beyond the apoptosis peak raises the question of whether they contribute to ongoing degeneration a possibility warranting further study using microglial depletion or inhibition strategies [12].

Our findings are broadly consistent with the existing literature on light-induced retinal degeneration [1-5, 15]. However, this study provides a detailed, side-by-side characterization of morphological, apoptotic, and inflammatory parameters within the critical first 72 hours, offering a comprehensive timeline in a single experimental system. The light-damage model has significant translational relevance. Age-related macular degeneration, the leading cause of elderly blindness, involves progressive photoreceptor degeneration and chronic inflammation [3, 6]. The acute events we documented photoreceptor apoptosis followed by microglial activation recapitulate key aspects of early AMD pathology [4]. Similarly, in retinitis pigmentosa, photoreceptor apoptosis is the final common pathway to vision loss, with inflammation increasingly recognized as a disease modifier [5]. The 24- to 48-hour window identified in our study represents an optimal therapeutic opportunity [16]. Anti-apoptotic strategies (caspase inhibitors, neurotrophic factors) would require administration before or shortly after insult [7]. Modulating the microglial response could be beneficial either by enhancing phagocytic clearance to promote recovery or dampening pro-inflammatory activity to limit bystander damage [14]. The challenge remains balancing these opposing functions for optimal retinal preservation.

Conclusion

In conclusion, this study demonstrates that within the first three days following light-induced retinal injury: (1) Photoreceptor apoptosis is the primary pathological event, peaking at 48 hours and declining by 72 hours; (2) Microglial cells transform from ramified to amoeboid morphology, proliferate, and migrate to the degenerating ONL, with peak accumulation between days two and three; (3) The peak

of microglial accumulation closely follows the peak of apoptosis, confirming a direct temporal relationship; and (4) The first 48 hours emerge as a critical period when photoreceptor death is most active and the inflammatory response is being established. These findings contribute to understanding initial pathophysiological processes in retinal degeneration and highlight a narrow therapeutic window. Future studies should focus on dissecting the molecular signals governing the interaction between apoptotic photoreceptors and responding microglia, with the ultimate goal of identifying targets for early therapeutic strategies aimed at preserving retinal structure and function.

Acknowledgments

The author would like to thank the Laboratory Animal Sciences Center for their support and assistance with animal housing and experimental procedures.

References

1. Organisciak DT, Vaughan DK. Retinal light damage: Mechanisms and protection. *Prog Retin Eye Res.* 2010;29(2):113-34.
2. Wenzel A, Grimm C, Samardzija M, Remé CE. Molecular mechanisms of light-induced photoreceptor apoptosis and neuroprotection for retinal degeneration. *Prog Retin Eye Res.* 2005;24(2):275-306.
3. Pennesi ME, Neuringer M, Courtney RJ. Animal models of age related macular degeneration. *Mol Aspects Med.* 2012;33(4):487-509.
4. Wright AF, Chakarova CF, Abd El-Aziz MM, Bhattacharya SS. Photoreceptor degeneration: genetic and mechanistic dissection of a complex trait. *Nat Rev Genet.* 2010;11(4):273-84.
5. Carter-Dawson LD, LaVail MM. Rods and cones in the mouse retina. I. Structural analysis using light and electron microscopy. *J Comp Neurol.* 1979;188(2):245-62.
6. Strauss O. The retinal pigment epithelium in visual function. *Physiol Rev.* 2005;85(3):845-81.
7. Hafezi F, Marti A, Munz K, Remé CE. Light-induced apoptosis: differential timing in the retina

- and pigment epithelium. *Exp Eye Res.* 1997;64(6):963-70.
8. Remé CE, Grimm C, Hafezi F, Marti A, Wenzel A. Apoptotic cell death in retinal degenerations. *Prog Retin Eye Res.* 1998;17(4):443-64.
9. Gavrieli Y, Sherman Y, Ben-Sasson SA. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol.* 1992;119(3):493-501.
10. Abler AS, Chang CJ, Ful J, Tso MOM, Lam TT. Photoc injury triggers apoptosis of photoreceptor cells. *Res Commun Mol Pathol Pharmacol.* 1996;92(2):177-189.
11. Karlstetter M, Scholz R, Rutar M, Wong WT, Provis JM, Langmann T. Retinal microglia: just bystander or target for therapy? *Prog Retin Eye Res.* 2015;45:30-57.
12. Silverman SM, Wong WT. Microglia in the retina: roles in development, maturity, and disease. *Annu Rev Vis Sci.* 2018;4:45-77.
13. Langmann T. Microglia activation in retinal degeneration. *J Leukoc Biol.* 2007;81(6):1345-51.
14. Santos AM, Calvente R, Tassi M, et al. Embryonic and postnatal development of microglial cells in the mouse retina. *J Comp Neurol.* 2008;506(2):224-39.
15. Noailles A, Fernández-Sánchez L, Lax P, Cuenca N. Microglia activation in a model of retinal degeneration and TUDCA neuroprotective effects. *J Neuroinflammation.* 2014;11:186.
16. Sachiko Saino-Saito & Reza Mohammad Nourani, Discrete localization of various fatty-acid-binding proteins in various cell populations of mouse retina. *Cell Tissue Res* (2009) 338:191–201
17. LaVail MM, Gorrin GM, Repaci MA, Thomas LA, Ginsberg HM. Genetic regulation of light damage to photoreceptors. *Invest Ophthalmol Vis Sci.* 1987;28(6):1043-8.
18. Hume DA, Perry VH, Gordon S. Immunohistochemical localization of a macrophage-specific antigen in developing mouse retina: phagocytosis of dying neurons and differentiation of microglial cells to form a regular array in the plexiform layers. *J Cell Biol.* 1983;97(1):253-7.
19. oammad Hamed Rashidi, Masoud Ghorbani, Mohammad Reza Nouran. Immunohistochemical Analysis of F4/80-Positive Macrophages in the Interstitial Tissue of Mouse Testis. *Life Sciences Student Journal*, 2024, vol2, No2 pp10-16
20. Aziz MK, Ni A, Quintana D, et al. Evidence of early ultrastructural photoreceptor abnormalities in light-induced retinal degeneration using spectral domain optical coherence tomography. *Br J Ophthalmol.* 2014;98(7):984-9.